


IN THE CLAIMS

1. (currently amended) A method for the synthesis of a KPV tripeptide diamide derivative represented by the following formula (I)



or for a salt thereof, independent of stereochemistry wherein:

- a) R_1 , R'_1 and R''_1 represent, independently from each other, a hydrogen atom or
- a linear or branched C_1 - C_{22} alkyl moiety, optionally interrupted by a heteroatom,
 - C_4 - C_{10} cycloalkyl moiety,
 - a linear or branched C_1 - C_{22} polyfluoroalkyl or perfluoroalkyl moiety,
 - an aryl moiety optionally substituted by one or more halogen atoms or more linear or branched C_1 - C_4 alkyl moieties,
 - an aralkyl moiety,
 - or R_1 and R'_1 could form with $C(R''_1)$ a saturated ring with from 3 to 7 atoms, optionally substituted by one or more linear or branched C_1 - C_4 alkyl moieties and/or optionally containing a heteroatom,

with the proviso that the $R_1(R'_1)(R''_1)CO$ group does not represent an amino acid residue or a peptide;

b) R_2 and R_3 represent, independently from each other, a hydrogen atom or represent

- a linear or branched C_1-C_{24} alkyl moiety, optionally interrupted by a heteroatom,

- a C_4-C_{10} cycloalkyl moiety,

- a linear or branched C_1-C_{22} polyfluoroalkyl or perfluoroalkyl moiety,

- an aryl moiety optionally substituted by one or more halogen atoms or one or more linear or branched C_1-C_4 alkyl moieties,


- an aralkyl moiety,

- or R_2 and R_3 could form with the nitrogen atom a saturated ring with from 5 or 6 atoms optionally substituted by one or more linear or branched C_1-C_4 alkyl moieties, said saturated ring optionally containing a heteroatom ,

with the proviso that the $N(R_2)(R_3)$ group does not represent an amino acid or a peptide;

said method comprising:


a) reacting a lysine diprotected residue having the following formula (II):



optionally salified by a mineral or organic base,


wherein P_1 and P_2 , are different and each represent independently from one another a protective group,

with a proline residue having the following formula (III):




optionally salified by a mineral or organic acid,
 wherein P_3 represents a protective group differing from any
 of the P_1 and P_2 protective groups, ~~or wherein P_3 represents a~~
~~hydroxyl group~~

in the presence of an activation reagent or a coupling
 reagent and in a solvent, so as to obtain the following compound
 having the formula (IV):





wherein P_1 , P_2 and P_3 have the above-mentioned meanings,
 b)


1) removing the P_1 protective group of the compound of
 formula (IV) to obtain the compound of formula (IX):





2) amidating the NH_2 (α) group of the lysine residue of the compound of the formula (IX) with a compound having formula (VI-A) or a compound having formula (VI-B) to obtain the compound of formula (X):



3) removing the P_3 protective group from the compound of formula (X) to obtain the compound of formula (XI):



4) coupling the compound of formula (XI) with the valine compound of formula (V), or the mineral or organic salt thereof, to form the compound of formula (XII):



c) removing the P₂ protective group from the compound of formula (XII) to obtain the compound of the formula (I) or the mineral or organic salt thereof.

2. (original) The method according to claim 1, wherein the compound having the formula (I) is a salt selected amongst the hydrochlorides, hydrobromides, sulphates, acetates, citrates, tartrates, lactates, phosphates, hydrogenophosphates, propionates and succinates.

3. (previously presented) The method according to claims 1 or 2, wherein the lysine, proline or valine amino acid residues are any of the stereoisomers of such residues.

4. (original) The method according to claims 1 or 2, wherein the salt is obtained during step c) through introducing the corresponding acid.

5. (original) The method according to claim 4, wherein the acid is acetic acid, hydrochloric acid, hydrobromic acid, sulphuric acid, citric acid, tartaric acid, lactic acid, phosphoric acid, hydrogenophosphoric acid, propionic acid or succinic acid.

6. (original) The method according to claim 5, wherein the acid is acetic or hydrochloric acid.

7. (previously presented) The method according to claims 1 or 2, wherein the P_1 and P_2 protective groups represent, independently from each other, Adoc (1-adamantylloxycarbonyl) BOC (t-butyloxycarbonyl), 2-bromo-Z (2-bromo-benzyloxycarbonyl), 2-chloro-Z (2-chloro-benzyloxycarbonyl), Fmoc (9-fluorenylmethoxycarbonyl), Formyl, Nicotinoyl, 4-nitro-Z (4-nitro-benzyloxycarbonyl), Tfa (trifluoroacetyl), Tos (p-toluenesulfonyl), Z (benzyloxycarbonyl) or Adpoc (1-(adamantyl)-1-methylethoxycarbonyl).

8. (previously presented) The method according to claims 1 or 2, wherein the P_1 and P_2 protective groups are selected such as to be removed respectively under distinct operating conditions.

9. (previously presented) The method according to claims 1 or 2, wherein the compound having the formula (II) is salified by an organic base.

10. (original) The method according to claims 1 or 2, wherein the compound having the formula (III) is salified by a mineral or an organic acid.

11. (previously presented) A method according to claims 1 or 2, wherein in step a), the peptide coupling reaction occurs in the presence of an activation or a coupling reagent selected amongst carbodiimides, water-soluble carbodiimides, phosphonium salts, PyBOP ((benzotriazol-1-yloxy)tripyrrolidino-phosphonium hexafluorophosphate), PyBROP (bromotripyrrolidino-

phosphonium hexafluorophosphate), PyCloP (chlorotripyrrolidino-phosphonium hexafluorophosphate), or also by means of reagents selected amongst PyClU (chloro-N,N,N',N'-bis(tetramethylene)formamidinium hexafluoro-phosphate), N-hydroxysuccinimide, EEDQ (1-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinolin), CDI (carbonyldiimidazole), or chloroformates

Claims 12 - 14 (cancelled)

15. (original) The method according to claims 1 or 2, wherein in the compound having the formula (II), the P₁ protective group is t-butyloxycarbonyl (BOC) and the P₂ protective group is benzyloxycarbonyl (Z).


16. (previously presented) The method according to claims 1 or 2, wherein in the compound of the formula (III), the P₃ protective group is the OBzl ester group.

17. (previously presented) The method according to claims 1 or 2, wherein in the compound having the formula (I), the R₁, R'₁ and R''₁ groups represent each a hydrogen atom.

18. (original) The method according to claims 1 or 2, wherein in the compound having the formula (I), the R₂ and R₃ groups represent each a hydrogen atom.

19. (previously presented) The method according to claims 1 or 2, wherein the P₁ protective group is t-butyloxycarbonyl (BOC), the P₂ protective group is benzyloxycarbonyl (Z) and the P₃ protective group is OBzl ester.

20. (withdrawn) A KPV tripeptide diamide derivative or salt thereof represented by the following formula (IA):



(IA)

wherein:

a) R₁, R'₁ and R'₁' represent, independently from each other, a hydrogen atom or

- a linear or branched C₁-C₂₂ alkyl moiety, optionally interrupted by a heteroatom,

- C₄-C₁₀ cycloalkyl moiety,

- a linear or branched C₁-C₂₂ polyfluoroalkyl or perfluoroalkyl moiety,

- an aryl moiety optionally substituted by one or more halogen atoms or one or more linear or branched C₁-C₄ alkyl moieties,

- an aralkyl moiety,

- or R₁ and R'₁ could form with C(R'₁) a saturated ring with from 3 to 7 atoms, optionally substituted by one or more linear or branched C₁-C₄ alkyl moieties and/or optionally containing a heteroatom,

- hydrogen,

with the proviso that the R₁(R'₁)(R'₁')CO group does not represent an amino acid residue or a peptide residue with at least one of R₁, R'₁, R'₁' being different from hydrogen.

b) R₂ and R₃ represent, independently from each other, a hydrogen atom or represent

- a linear or branched C_1-C_{24} alkyl moiety, optionally interrupted by a heteroatom,
 - a C_4-C_{10} cycloalkyl moiety,
 - a linear or branched C_1-C_{22} polyfluoroalkyl or perfluoroalkyl moiety,
 - an aryl moiety optionally substituted by one or more halogen atoms, or one or more linear or branched C_1-C_4 alkyl moieties,
 - an aralkyl moiety,
 - or R_2 and R_3 could form with the nitrogen atom a saturated ring with from 5 or 6 atoms optionally substituted by one or more linear or branched C_1-C_4 alkyl moieties, said saturated ring optionally containing a heteroatom, with at least one of the residues R_2 or R_3 being different from hydrogen,
- with the proviso that the $N(R_2)$ (R_3) group does not represent an amino acid or a peptide residue.

21. (withdrawn) The KPV tripeptide diamide derivative according to claim 20, wherein the salt is selected amongst hydrochlorides, hydrobromides, sulphates, acetates, citrates, tartrates, lactates, phosphates, hydrogenophosphates, propionates and succinates.

22. (withdrawn) The KPV tripeptide diamide derivate according to claims 20 or 21, wherein the Lysine, Proline or Valine amino acid residues are any of the stereoisomers of each of such residues.

23. (withdrawn) A composition comprising: a KPV tripeptide diamide derivative or salt thereof according to claims 20 or 21 in a physiologically acceptable medium.

24. (withdrawn) The composition according to claim 23, wherein the physiologically acceptable medium is a cosmetic medium and the KPV tripeptide diamide derivate or salt thereof is present in an amount ranging from 10^{-8} to 10^{-3} g/100g.

25. (withdrawn) The composition according to claim 23, wherein the physiologically acceptable medium is a pharmaceutical medium and the KPV tripeptide diamide derivate is present in an amount greater than $5 \cdot 10^{-4}$ g/100g.

26. (cancelled)

27. (original) The method according to claim 9, wherein the organic base is an organic amine.

28. (cancelled)

29. (withdrawn) A method of treating dry or sensitive skin comprising: obtaining a quantity of a composition of claim 23 and applying said composition to the dry or sensitive skin of a patient.

30. (withdrawn) A method of treating dry or sensitive skin comprising: obtaining a quantity of a composition of claim 24 and applying said composition to the dry or sensitive skin of a patient.

31. (withdrawn) A method of treating dry or sensitive skin comprising: obtaining a quantity of a composition of claim 25 and applying said composition to the dry or sensitive skin of a patient.

32. (withdrawn) The method of claim 1 wherein R_1 , R'_1 or R''_1 are a linear or branched C_1 - C_{22} alkyl moiety interrupted by a heteroatom, said heteroatom is selected from O, N, S or Si.

33. (withdrawn) The method of claim 1 wherein when R_1 and R'_1 form with $C(R''_1)$ a saturated ring containing a heteroatom, said heteroatom is O, S or N.

34. (withdrawn) The method of claim 1 wherein when R_2 and R_3 is a linear or branched C_1 - C_{22} alkyl moiety interrupted by a heteroatom, said heteroatom is selected from O, N, S or Si.

35. (withdrawn) The method of claim 1 wherein when R_2 and R_3 form with a nitrogen atom a saturated ring containing a heteroatom, said heteroatom is O, S or N.

36. (withdrawn) The method of claim 9 wherein said organic base is an organic amine.

37. (withdrawn) The method of claim 1 wherein when R_1 , R'_1 or R''_1 are an aryl moiety optionally substituted by one or more halogen atoms, such halogen is Cl, F, Br or I.

38. (withdrawn) The method of claim 1 wherein when R_2 and R_3 form an aryl moiety optionally substituted by one or more halogen atoms, such halogen is Cl, F, Br or I.

39. (withdrawn) The KPV tripeptide diamide derivative or salt thereof of claim 20 wherein, when R_1 , R'_1 and R''_1 represent a linear or branched C_1 - C_{22} alkyl moiety interrupted by a heteroatom, said heteroatom is O, N, S or Si.

40. (withdrawn) The KPV tripeptide diamide derivative or salt thereof of claim 20 wherein, when R_1 , R'_1 and R''_1 represent an aryl moiety optionally substituted by one or more halogen atoms, such halogen is Cl, F, Br or I.

41. (withdrawn) The KPV tripeptide diamide derivative or salt thereof of claim 20 wherein, when R_1 and R'_1 form with $C(R''_1)$ a saturated ring containing a heteroatom, said heteroatom is O, S or N.

42. (withdrawn) The KPV tripeptide diamide derivative or salt thereof of claim 20 wherein, when R_2 and R_3 represent a linear or branched C_1 - C_{22} alkyl moiety interrupted by a heteroatom, said heteroatom is O, N, S or Si.

43. (withdrawn) The KPV tripeptide diamide derivative or salt thereof of claim 20 wherein, when R_2 and R_3 represent an aryl moiety optionally substituted by one or more halogen atoms, such halogen is Cl, F, Br or I.

44. (withdrawn) The KPV tripeptide diamide derivative or salt thereof of claim 20 wherein, when R_2 and R_3 form with a nitrogen atom a saturated ring containing a heteroatom, said heteroatom is O, S or N.

45. (withdrawn) A method of making a composition useful for treating dry or sensitive skin comprising obtaining a quantity of a KPV tripeptide diamide derivative or salt thereof as claimed in claim 20 and mixing same in a physiologically acceptable medium so as to produce a dermatological composition.

46. (previously presented) The method of claim 1, which does not comprise a final purification step.